



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

We thus arrive at the result:

$$\binom{cn}{a} - \binom{a}{1} \binom{cn-c}{a} + \binom{a}{2} \binom{cn-2c}{a} - \cdots + (-1)^a \binom{cn-ca}{a} = c^a.$$

## II. SOLUTION BY NORMAN ANNING, Chilliwack, B. C.

Define operators  $E$  and  $\Delta$  as follows:

$$E^h f(n) = f(n+h), \quad \Delta f(n) = f(n+1) - f(n) = (E-1)f(n).$$

Then the given expression,

$$\begin{aligned} \binom{cn}{a} - \binom{a}{1} \binom{cn-c}{a} + \binom{a}{2} \binom{cn-2c}{a} - \cdots + (-1)^a \binom{cn-ca}{a} \\ = (1 - E^{-1})^a \binom{cn}{a} = (\Delta E^{-1})^a \binom{cn}{a} = \Delta^a E^{-a} \binom{cn}{a} = \Delta^a \binom{cn-ca}{a}. \end{aligned}$$

Now

$$\binom{cn-ca}{a} = \frac{1}{a!} \{c^a n^a + \text{terms in } n \text{ of degree lower than } a\}.$$

Hence,

$$\Delta^a \binom{cn-ca}{a} = \frac{1}{a!} \{c^a \cdot a! + 0\} = c^a.$$

### 446. Proposed by CLIFFORD N. MILLS, Brookings, South Dakota.

Solve the equations

$$x^2(y-z) = l^2(m-n), \quad y^2(z-x) = m^2(n-l), \quad z^2(x-y) = n^2(l-m).$$

## SOLUTION BY H. S. UHLER, Yale University.

An equation involving only one variable  $x$  may be obtained in the following manner. Equating the expressions for  $z$  derived from the first two given equations we find

$$y - l^2(m-n)/x^2 = x + m^2(n-l)/y^2$$

or

$$x^2 y^2 - x^2 y^2 + m^2(n-l)x^2 + l^2(m-n)y^2 = 0. \quad (1)$$

Substituting the value of  $z$  from the first in the third given equation and reducing, we obtain

$$x^5 y^2 - x^4 y^2 - n^2(l-m)x^4 - 2l^2(m-n)x^2 y + 2l^2(m-n)x^2 y^2 + l^4(m-n)^2 x - l^4(m-n)^2 y = 0. \quad (2)$$

Multiplying (1) by  $x^2$  and subtracting (2) from the product gives, after removing the factor  $m-n$ ,

$$(lm-mn+nl)x^4 - 2l^2 x^2 y + l^2 x^2 y^2 + l^4(m-n)x - l^4(m-n)y = 0. \quad (3)$$

Multiplying (1) by  $l^2$ , (3) by  $y$ , adding, and dividing by  $x$ ,

$$(lm-mn+nl)x^2 y - l^2 x^2 y^2 + l^2 m^2(n-l)x + l^4(m-n)y = 0. \quad (4)$$

Adding (3) and (4) and dividing by  $x$ ,

$$(lm-mn+nl)x^3 + (lm-mn+nl-2l^2)x^2 y + l^2[l^2(m-n) + m^2(n-l)] = 0. \quad (5)$$

Substituting the expression for  $y$  from (5) in (4) and removing the factors  $(l-m)(l-n)$  we derive the following quadratic in  $x^2$ , namely

$$(lm-mn+nl)^2 x^6 - 2l^4[m^2(l-n) + n^2(l-m)]x^3 - l^6(mn-nl+lm)(nl-lm+mn) = 0.$$

Consequently

$$x^3 = l^3$$

and

$$x^3 = \frac{l^3[l^2(m-n)^2 - m^2 n^2]}{(lm-mn+nl)^2}.$$

Finally, the six values of  $x$  are

$$l, l\omega, l\omega^2, \frac{l[l^2(m-n)^2 - m^2n^2]^{\frac{1}{2}}}{(lm - mn + nl)^{\frac{2}{3}}}, \frac{l[l^2(m-n)^2 - m^2n^2]^{\frac{1}{2}}\omega}{(lm - mn + nl)^{\frac{2}{3}}}, \frac{l[l^2(m-n)^2 - m^2n^2]^{\frac{1}{2}}\omega^2}{(lm - mn + nl)^{\frac{2}{3}}},$$

where

$$\omega = (-1 + i\sqrt{3})/2.$$

The corresponding values of  $y$  and  $z$  may be obtained at once by cyclical permutation of the letters  $l, m$ , and  $n$ .

*Remarks:* The above analysis was verified by direct substitution of the six sets of values in the given equations. The elimination of  $y$  between equations (1) and (2) can be performed in a more elegant manner by Sylvester's dialytic method, but the simplification of the determinant of the eighth order requires too much space for publication.

Also solved by A. H. HOLMES and NORMAN ANNING.

#### GEOMETRY.

##### 471. Proposed by C. N. SCHMALL, New York City.

In the ellipse  $x^2/a^2 + y^2/b^2 = 1$ , an equilateral hexagon is inscribed with two sides parallel to the major axis. In the major auxiliary circle the same thing is done. If  $H_1$  and  $H_2$  be the sides of the hexagons, and  $e$  the eccentricity of the ellipse, show that  $H_1 : H_2 :: 4 - 2e^2 : 4 - e^2$ .


SOLUTION BY J. A. CAPARO, University of Notre Dame.

Since  $AD$  is the side of a regular hexagon inscribed in the major circle of the ellipse  $x^2/a^2 + y^2/b^2 = 1$ , we have

$$DA = OA = a = H_2$$

and since

$$e^2 = \frac{a^2 - b^2}{a^2}, \quad b^2 = H_2^2(1 - e^2), \quad \therefore \frac{x^2}{H_2^2} + \frac{y^2}{H_2^2(1 - e^2)} = 1. \quad (1)$$



Now  $EF = H_1/2$ ; therefore, from (1) we have

$$FC^2 = y^2 = \frac{(4H_2^2 - H_1^2)(1 - e^2)}{4}.$$

Also  $CA = OA - OC$ . Hence,  $CA = H_2 - (H_1/2)$ ; and since  $FA^2 = FC^2 + CA^2$ , we have

$$4H_1^2 = (4H_2^2 - H_1^2)(1 - e^2) + (2H_2 - H_1)^2.$$

Write  $H_1/H_2 = x$ . Then, substituting and reducing, we have

$$x^2(4 - e^2) + 4x + 4(e^2 - 2) = 0.$$

Solving for  $x$ , we have

$$x = \frac{-2 \pm 2(e^2 - 3)}{4 - e^2}.$$

Using the negative sign, we have  $x = (4 - 2e^2)/(4 - e^2)$ ; and since  $x = H_1/H_2$ ,

$$H_1 : H_2 :: 4 - 2e^2 : 4 - e^2.$$

Also solved by C. E. HORNE, FRANK IRWIN, HAROLD T. DAVIS, NORMAN ANNING, ELIJAH SWIFT, and C. N. SCHMALL.